arXiv:cond-mat/0703290v1 [cond-mat.supr-con] 12 Mar 2007

Observed Non-Fermi Liquid Heat Transport and Entropy of Pseudogap Quasi-Particles as Possible Manifestations of Single Particle Bosons

B. Abdullaev, C. -H. Park, and K. -S. Park
Research Center for Dielectric and Advanced Matter Physics,
Department of Physics, Pusan National University,
30 Jangjeon-dong, Geumjeong-gu, Busan 609-735, Korea.

(Dated: Received February 6, 2008)

We show that the ideal gas of Bogolyubov quasi-particles (IGBQ) of $2D$ Coulomb Bose gas provides $c \sim T^4$ and $c \sim T$ (from plasmon and free particle parts of quasi-particles, respectively) temperature dependencies of the specific heat. This can in a framework of Drude model interpret the observed low-temperature heat conductivity $\kappa \sim T^{3.6}$ and $\kappa \sim T$ of pseudogap quasi-particles of cuprate superconductors. Vanishing of free quasi-particles of IGBQ in the transition to plasmons determines the downturn behavior of κ and clarifies the origin of decoupling of charge carriers with heat. Near to pseudogap boundary Bose gas transforms into Fermi gas of holes or electrons. Obtained from mixture of Bose and Fermi gases Lorentz ratio values of Wiedemann – Franz law (WFL) are close to observed non-Fermi liquid ones. The observed temperature dependence of entropy can be result of the specific heat contributions from these both gases.

PACS numbers: 74.72.-h, 74.20.Mn, 74.25.Fy, 74.25.Bt

In the seminal paper [1] on investigation the low-temperature WFL for heat transport, Hill with collaborators have experimentally studied the properties of normal state quasi-particles in the electron doped $Pr_{2-x}Ce_xCuO_4$ copper-oxide. It was shown by Hill et al. that they failed the Landau Fermi liquid theory (LFLT). In the high external magnetic field the superconductivity was suppressed and the heat conductivity of quasi-particles had a linear $\kappa \sim T$ law at a decreasing the temperature up to some value and then undergone $\kappa \sim T^{3.6}$ downturn behavior. The latter was called in the literature a downturn of heat conductivity. Next important result of [1] was Lorentz ratio did not equate to Sommerfeld's value. These two observations were regarded as evidences of breakdown of LFLT in cuprate superconductors.

The downturn was observed in Ref. [2] and in the heavily over-doped non-superconducting compound $La_{2-x}Sr_xCuO_4$ at low temperatures. The violation of WFL in the normal state of $Bi_{2+x}Sr_{2-x}CuO_{6+\delta}$ copper-oxide in the vicinity of metal-insulator transition was also detected in the recent paper [3].

Now it is widely believed [3] that the phenomenological explanation of downturn is in the decoupling of phonon thermal bath with current producing charge carriers [4] and therefore, has no relation to fundamental properties of high-T_c superconductors. In contrast, Hill et al. have supposed that the downturn may be intrinsic for copper-oxides. It would be interesting to find the experimental revealing of the downturn effect of heat conductivity in the another observable quantity, if it would be intrinsic, and understand the general underlying physics.

It seems the experiment gives us this opportunity. Namely, no T dependence observation [2, 3] of the electric conductivity at low-temperatures allows to explicitly connect in the framework of Drude model the temperature dependence of heat conductivity κ with one of the specific heat c.

The existing observed results on c of cuprate materials are as follows. The normal state electronic specific heat of high-T_c superconductors $YBa_2Cu_3O_{6+x}$ and $La_{2-x}Sr_xCuO_4$ was experimentally investigated in [5] and [6], respectively. Its magnetic field dependence of $Y_{0.8}Ca_{0.2}Ba_2Cu_3O_{6+x}$ compound has considered in [7]. The existence of superconductivity at low temperatures does not allow to extract the T dependence of the normal state c. However, the normal state properties are clearly seen [5] in the entropy $\mathcal{S}$. For instance, the doping dependence of $\mathcal{S}$ in the $YBa_2Cu_3O_{6+x}$ coper-oxide was displayed in [5]. It was also shown in [5] that for under-doped material the low-temperature dependence was $\mathcal{S} \sim T^n$ with $n > 1$, while $\mathcal{S} \sim T$ was obtained for optimal doping compound. Non-linear T dependence of the entropy was related to isolating state [5] of under-doped materials.

One might assume that $\kappa \sim T^{3.6}$ and $\mathcal{S} \sim T^n$ have the same underlying physical origin and could be expressed through T dependence of the common specific heat c. In this case downturn of κ would have a physical meaning and correspond to isolating state, which is consistent with the decoupling mechanism [4] of thermal phonons with current carrying particles.

In the present paper we follow Ref. [8] that high-T_c superconductivity (HTCS) and pseudogap (PG) regime are described by mechanism of the single particle $2D$ Coulomb interacting bosons, in which the normal state of superconductors is IGBQ. The physical origin of bosons is in the bosonization of $2D$ fermions below PG boundary due to coupling of their spins with statistical (anyon) magnetic field. According this mechanism in the deeply under-doped regime and low temperatures dominate bosons, while close to PG boundary there occurs the decoupling of spins and statistical magnetic field and dominate fermions. Therefore, the normal state is mixture of bosons and fermions. In [8] we have used the Bogolyubov approximation for $2D$ Coulomb Bose gas, which is valid only in the high density limit of gas. Being applied to gas from neutral atoms, this approximation has validity for weakly dilute densities. However, as it was shown in Ref. [9], it can give good results also for superfluid He II. Thus, we can assume that our results for bosons are applicable for intermediate densities of $2D$ Coulomb Bose gas.

We do the calculation of the specific heat of IGBQ. For ideal Bose gas the thermodynamic free energy has the form [10]:

$$F = T\int \frac{d^2\,p\,d^2\,x}{(2\pi\hbar)^2}\,Ln(1-e^{-\varepsilon(p)/T})\,. \qquad (1)$$

The energy scale is implied for temperature T. In the energy of quasi-particles [8] $\varepsilon(p) = (ap + (p^2/2m)^2)^{1/2}$ ap is the square of plasmon energy with $a = 2\pi e^2 n\hbar/m$, where e and n are the charge and $2D$ density, respectively, of true particles with mass m, and p is momentum of quasi-particles.

The integral in Eq. (1) over the absolute value of momentum p we write in the form: $\int_0^\infty = \int_0^q + \int_q^\infty$. Here q provides at $T = 0$ the maximum of the momentum distribution function for true non-condensate particles [11] and separates the plasmon and kinetic energy contributions to energy of quasi-particles.

If for the first (second) integral we introduce the variable $x = (ap)^{1/2}/T$ ($x = p^2/(2mT)$) and then take into account that $(aq)^{1/2} = q^2/(2m)$ we obtain

$$F \;\approx\; \frac{TS}{2\pi\hbar^2}\left[\frac{2T^4}{a^2}\int\limits_0^{(aq)^{1/2}/T} d\,x\,x^3\,Ln(1-e^{-x}) \right.$$
$$\left. +\;\; m\,T\int\limits_{(aq)^{1/2}/T}^{\infty} d\,x\,Ln(1-e^{-x})\right]\,. \qquad (2)$$

Here S is the area of $2D$ system.

For the low temperatures, when $(aq)^{1/2}/T \gg 1$, we can replace the upper limit of first integral in Eq. (2) by infinity sign. In this case the second integral becomes zero. For the high temperatures we have $(aq)^{1/2}/T \ll 1$ then the lower limit of second integral can be putted zero and zero becomes the first integral.

Following the scheme of calculation [10], we find the expressions for the specific heat divided to S

$$c_1 = -\frac{20T^4A}{\pi\hbar^2 a^2} \qquad (3)$$

for ideal gas of plasmons and

$$c_2 = -\frac{TmB}{\pi\hbar^2} \qquad (4)$$

for ideal gas of free quasi-particles. Here $A = -(1/4)\int_0^\infty x^4dx/(e^x-1) = -(1/4)\Gamma(5)\zeta(5)$ and $B = -\int_0^\infty xdx/(e^x-1) = -\Gamma(2)\zeta(2)$, where $\Gamma(y)$ and $\zeta(y)$ are Gamma and the Riemann Zeta functions, respectively.

The normal state heat conductivity κ is described by WFL for free quasi-particles (charged bosons). Its charge conductivity σ, at low-temperatures, can approximately be obtained from the Drude model. The gas of plasmons does not care the charge therefore, is isolating. However, to determine the approximate downturn temperature T_d we can formally extend the WFL to the region of this isolating gas. The transition from electrically conducting into non-conducting case will be described as transition from finite to zero (as it should be for isolators) charge conductivity.

So, the estimate of downturn temperature T_d is determined from equation

$$\kappa_1(T_d)/\sigma_1 = \kappa_2(T_d)/\sigma_2\,. \qquad (5)$$

Here the Drude model gives the expressions of heat conductivities for gas of plasmons $\kappa_1(T_d) = (1/2)c_1(T_d)v_1l_1$ and gas of free quasi-particles $\kappa_2(T_d) = (1/2)c_2(T_d)v_2l_2$. Corresponding mean velocities, free penetration lengths and charge conductivities are $v_{1,2}$, $l_{1,2}$ and $\sigma_{1,2}$, respectively.

The solution of equation Eq. (5) is

$$T_d = \left(\frac{\Gamma(2)\zeta(2)}{5\Gamma(5)\zeta(5)}\right)^{1/3}\left(\frac{\sigma_1 v_2 l_2}{\sigma_2 v_1 l_1}\right)^{1/3}(ma^2)^{1/3}\,. \qquad (6)$$

One can show that $(ma^2)^{1/3} = 2(2/r_s^2)^{2/3}Ry$, where r_s is the mean distance between particles in Bohr radius, a_B ($a_B = \hbar^2/(me^2)$), units and $Ry = me^4/(2\hbar^2)$ is Rydberg energy.

We use the density of charges [8] $n_{ab} = 6.747\cdot 10^{14}\cdot t\cdot 1/cm^2$ per $1\,cm^2$ area with doping fraction t of charge per Cu atom. It is almost the same for all cuprates, which is result of equivalent spacing constants [12] for elementary structural cells of CuO_2 current carrying $a-b$ planes. We rewrite this density through a_B units then r_s will be the function of t. T_d has a form:

$$T_d \approx 0.114\left(\frac{\sigma_1 v_2 l_2}{\sigma_2 v_1 l_1}\right)^{1/3} t^{2/3}Ry\,. \qquad (7)$$

Eq. (7) is an estimate of T_d. We do the numerical calculation of the specific heat for arbitrary T to precisely obtain T_d.

The expression for the specific heat is

$$c = \frac{q^2}{2m}\frac{m}{4\pi\hbar^2}\int_0^\infty \frac{d\,y}{z^2}\frac{y^2(1+y^3)}{sinh^2(y^{1/2}(1+y^3)^{1/2}/(2z))}\,, \quad (8)$$



FIG. 1: The specific heat c/z (in $q^2/(8\pi\hbar^2)$ units) as function of z.

where $z = T/(aq)^{1/2}$ and $y = p/q$. The result of numerical calculation of c/z (in $q^2/(8\pi\hbar^2)$ units) as function of z is plotted in Fig. 1. This quantity equals to $(4\pi\hbar^2/m)c/T$. In the assumption that heat conductivity is described by Drude model, $\kappa = mv^2c\sigma/(2ne^2)$, and σ does not depend from T (as was observed in [2, 3]), it displays temperature dependence of κ/T (we note that for degenerate Fermi liquid and $2D$ Coulomb Bose gas v is function only of concentration n). Experiments [2, 3] provide unique opportunity when only the specific heat c describes T dependence of κ and we can compare this result for κ with one for entropy $\mathcal{S}$.

From Fig. 1 we see that downturn of c/T occurs at $(aq)^{1/2}/2$. Calculation gives $(aq)^{1/2} = (16\pi e^2 n\hbar/(3(2m)^{1/2}))^{2/3} = 8(2^{1/2}/(3r_s^2))^{2/3}Ry$. Expressing r_s through t one obtains $(aq)^{1/2} = 0.737 \cdot t^{2/3}Ry$ and exact downturn temperature T_d, Eq. (7), with numerical factor 0.368.

We estimate quantities at $t \sim 0.1$. From expression for $(aq)^{1/2}$ one derives $(aq)^{1/2}/k_B \sim 10^4\,K$, therefore, the downturn temperature of the specific heat takes place at $10^4\,K$. Observed downturn of heat conductivity [1, 3] is $T_{d,exp}/k_B \sim 0.1\,K$. Using Eq. (7) we find $\sigma_1 v_2 l_2/(\sigma_2 v_1 l_1) \sim 10^{-12}$. One can assume $v_1 l_1 \sim v_2 l_2$, i.e. the product of mean velocities and free penetration lengths for plasmons and free quasi-particles has the same order of magnitude. Hence, we obtain $\sigma_1/\sigma_2 \sim 10^{-12}$. This result reflects the above supposed transition from charge conducting gas from free quasi-particles to isolating one from plasmons, i.e. metal-isolator transition. Therefore, the downturn of heat conductivity might be result of metal-isolator transition. It is worth here to note that the infinitesimal ratio of charge conductivities maps large scale of the specific heat temperatures of IGBQ into low ones of heat transport. We will discuss the metal-isolator crossover of cuprates within the single particle boson mechanism of HTCS elsewhere.

We calculate the Lorentz ratio for free quasi-particles of IGBQ. The WFL, $\kappa/(\sigma T) = L_0$, of heat transport for quasi-particles of LFLT is expressed through Lorentz ratio L_0 by Sommerfeld's value $L_0 = (\pi^2/3)(k_B/e)^2$. We note that this value of L_0 corresponds to the three dimensional ($3D$) case. A simple calculation shows that it does not change and for $2D$. For the IGBQ we assume that the mean kinetic energy of quasi-particles is $mv^2/2 = (aq)^{1/2}$ (because as was pointed out above at $T = 0$ the most part of non-condensate particles has this energy). Then the WFL, $\kappa/(\sigma T) = mv^2c/(2ne^2T)$, with c determined from Eq. (4) for free quasi-particles of IGBQ has a form $\kappa/(\sigma T) = 3.106 \cdot L_0/t^{1/3}$. This expression corresponds to WFL of bosons when there is no mixture of bosons with Fermi quasi-particles of LFLT. However, at the increasing of concentrations of holes or electrons t to the direction of PG boundary there appear the fermions in the PG region [8] and this mixture occurs. Therefore, at $T \to 0$ we can phenomenologically write the expression $\kappa/(\sigma T) = L$ with

$$L = L_0\left[\frac{3.106}{t^{1/3}}\left(1 - \frac{t}{t_c}\right) + \frac{t}{t_c}\right], \tag{9}$$

which takes into account the transformation of L from one of free quasi-particles of IGBQ to one of LFLT quasi-particles when concentration t tends to (but below) critical concentration $t_c \approx 0.19$ [13].

We do a comparison of our L/L_0 with the experimental one of [3]. In Fig. 7 (a) of [3] the data of L/L_0 for several copper-oxides are plotted as function of t. This dependence of t is almost universal for all cuprates. We use results for $Bi_{2+x}Sr_{2-x}CuO_{6+\delta}$ compound and obtain for t: 0.143, 0.17 and 0.185 the theoretical values of $(L/L_0)_{theor}$: 2.222, 1.485 and 1.117, while the experimental ones of $(L/L_0)_{exp}$ are 3.1 ± 0.7, 1.2 ± 0.2 and 1.05 ± 0.1, respectively.

Now before discussion of our result on entropy, we would like to outline the some observed qualitative features [5, 6, 13] of the HTCS specific heat and entropy. They are as follows. The increment coefficient $\gamma = c/T$ of the specific heat in the HTCS-to-normal-state phase transition (PT) point has weakly apparent and washed out peak in the deeply under-doped site. The peak of γ and its form become higher and sharper, respectively, when increases t. Peak is maximal (with the sharpest form) at t_c and then does not almost change the form for $t > t_c$. There is no influence [7] of the external magnetic field on the normal state c. It is observed the loss [7] of the normal state entropy $\mathcal{S}$ in cuprate when doping changes from optimal doping to under-doping, which is a result [5] of changing of $\mathcal{S} \sim T$ into $\mathcal{S} \sim T^n$ with $n > 1$. And last observation is γ does not depend from doping for $t > t_c$.

In general, the weakly apparent peak of γ is attributed [10] to first order PT, which is close to second order one. Sharpening of γ at t_c might characterize the transition of PT into second order type, where conventional superconductivity scenario with Cooper pairs becomes actual. This hypothesis was pointed out in [8].

No dependence of the normal state c from the magnetic field might be result that quasi-particle energy $\varepsilon(p)$ of IGBQ is independent of magnetic field. To outline it we follow the argument of Abrikosov, given in the book [14], for Cooper pair quasi-particle energy $\varepsilon_{Cp}(p)$, which is independent of magnetic field. He has shown that $\varepsilon_{Cp}(p)$, which is result of canonical Bogolyubov transformation, is function of transformational scalar quantities u_p and v_p. The latter ones in the magnetic field with gauge $\vec{\nabla}\cdot\vec{A}(\vec{r}) = 0$ can be function of only zero scalar product $\vec{k}\cdot\vec{A}_{\vec{k}} = 0$, where $\vec{k}$ and $\vec{A}_{\vec{k}}$ are Fourier transforms of vector-coordinate $\vec{r}$ and vector-potential $\vec{A}(\vec{r})$, respectively.

The calculation of the specific heat c_F of $2D$ gas of LFLT quasi-particles gives the same linear T dependence as in Eq. (4), but with ratio $c_F/c_2 = 2$. This is due to two directions of spins for fermions instead of bosons. From Eq. (4) it is seen that c_F is concentration independent. Therefore, observed no t dependence of γ for over optimal doping of copper-oxides proves the $2D$ nature of cuprates.

We find the expression of the normal state entropy $\mathcal{S}$. It is obtained through the calculation of integral $\mathcal{S} = \int_0^T (c/T_1) dT_1$. Our goal is to consider again the PG region, thus $T \leq T^*$, where T^* is PG boundary temperature. Typical experimental scale of T^* is $T^*/k_B \sim 10^2\, K$, while $(aq)^{1/2}/k_B \sim 10^4\, K$, therefore, in Eq. (8) one can assume $z \ll 1$ and use Eq. (3) as approximate expression of the specific heat of IGBQ. For arbitrary t and T (in the intervals of $t < t_c$ and $T \leq T^*$) we write the phenomenological expression

$$\frac{c}{T} = \frac{c_1}{T}\left(1 - \frac{5T}{4T^*}\right) + \frac{c_F}{T}\frac{T}{T^*}\,. \tag{10}$$

Except T/T^* both terms of this formula should also been include the term $\pi\xi^2(t)/(\pi\xi^2(t_c))$ with the same sign and numerical factors as for T/T^*. Here $\pi\xi^2(t)$ is area [8] of effective spin fluctuations of bosons at concentration t and $T = 0$, where bosons reveal fermion property and their spins interact with spins of Anti-Ferromagnetic environment atoms of cuprates. From Fig. 4 of paper [8] we find that $\pi\xi^2(t \to t_c) \to \infty$ and $\pi\xi^2(t < t_c)/(\pi\xi^2(t_c)) \approx 0$. Therefore, $\pi\xi^2(t)/(\pi\xi^2(t_c))$ term can be omitted in Eq. (10). The factor 5/4 in front of first T/T^* term was introduced for convenience purpose. The integration over T_1 gives

$$\mathcal{S} = \frac{c_1}{4}\left(1 - \frac{T}{T^*}\right) + \frac{c_F}{2}\frac{T}{T^*} \tag{11}$$

with qualitative experimental temperature dependencies of $\mathcal{S}$ in under-doping and near optimal doping.

In the Eqs. (9) and (11) we have used by hand introduced linear power dependencies on t/t_c and T/T^*, respectively. Further microscopical calculation can establish their exact law. However, the goal of the present paper was to show that at small these parameters dominant becomes the single particle boson contribution into low temperature heat conductivity and entropy, which is probably consistent with experiment.

As summary we have shown that observed normal state low-temperature dependencies of the heat conductivity $\kappa \sim T^{3.6}$ and $\kappa \sim T$ and entropy $\mathcal{S} \sim T^n$ with $n > 1$ for under-doped and $\mathcal{S} \sim T$ for optimal doped cuprate superconductors could be the result of functions $c \sim T^4$ and $c \sim T$ of the specific heat c. The temperature dependencies of c were obtained for mixture of IGBQ of $2D$ Coulomb Bose gas and gas of $2D$ LFLT quasi-particles. The obtained from this mixture Lorentz ratio values of WFL were close to observed non-Fermi liquid ones. The vanishing of free quasi-particles of IGBQ in the transition to plasmons has microscopically clarified the decoupling of charge carriers with heat and thus, the downturn of heat conductivity κ.

Acknowledgement. Author B. A. gratefully acknowledges the Korean Research Foundation Grant (KRF-2004-005-C00044) for support of work.

[1] R. W. Hill, C. Proust, L. Taillefer, P. Fournier, and R. L. Greene, Nature **414**, 711 (2001).

[2] S. Nakamae, K. Behnia, N. Mangkorntong, M. Nohara, H. Takagi, S. J. C. Yates, and N. E. Hussey, Phys. Rev. B **68**, 100502(R) (2003).

[3] C. Proust, K. Behnia, R. Bel, D. Mande, and S. I. Vedeneev, Phys. Rev. B **72**, 214511 (2005); cond-mat/0505551.

[4] M. F. Smith, J. Paglione, M. B. Walker, and L. Taillefer, Phys. Rev. B **71**, 014506 (2005).

[5] J. W. Loram, K. A. Mirza, J. R. Cooper, and W. Y. Liang, Phys. Rev. Lett. **71**, 1740 (1993).

[6] J. W. Loram, J. Luo, J. Cooper, W. Liang, and J. Tallon, J. Phys. Chem. Solids, **62**, 59 (2001).

[7] J. L. Luo, J. W. Loram, T. Xiang, J. R. Cooper, and J. L. Tallon, cond-mat/0112065.

[8] B. Abdullaev, in *Trends in Boson Research*, Editor A. V. Ling (Nova Science Publishers, N. Y., 2006), 139, cond-mat/0507500; B. Abdullaev,and C.-H. Park, J. Korean Phys. Soc. **49**, S642 (2006), cond-mat/0404648.

[9] N. M. March, W. H. Young, and S. Sampanthar, *The Many Body Problems in Quantum Mechanics* (Univ. Press, Cambridge, 1967).

[10] L. D. Landau, and E. M. Lifshitz, *Statistical Physics, Part 1* , (Pergamon Press, Oxford, 1980).

[11] E. M. Lifshitz, and L. P. Pitaevskii, *Statistical Physics, Part 2* , (Pergamon Press, Oxford, 1980).

[12] E. Dagotto, Rev. Mod. Phys. **66**, 763 (1994).

[13] J. L. Tallon, and J. W. Loram, Physica C **349**, 53 (2001).

[14] A. A. Abrikosov, *Fundamentals of the Theory of Metals*, (Elsevier Science, 1988).